

(19)  **Europäisches Patentamt**
European Patent Office
Office européen des brevets



(11) **EP 1 405 572 A1**

(12) **EUROPEAN PATENT APPLICATION**

(43) Date of publication: **07.04.2004 Bulletin 2004/15**
(51) Int Cl.7: **A23L 1/303, A23J 3/34, A23L 1/275, A23L 1/30**
(21) Application number: **02022479.6**
(22) Date of filing: **04.10.2002**

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR IE IT LI LU MC NL PT SE SK TR
Designated Extension States:
AL LT LV MK RO SI
(71) Applicant: **Fraunhofer-Gesellschaft zur Förderung der angewandten Forschung e.V.**
80686 München (DE)
(72) Inventors:
• **Funda, Elger**
79639 Grenzach-Wyhlen (DE)

• **Leuenberger, Bruno**
4123 Allschwil (CH)
• **Beck, Markus**
79540 Lörrach (DE)
• **Schäfer, Christian**
85368 Wang-Hagsdorf (DE)
(74) Representative: **Olgemöller, Luitgard, Dr. et al**
Leonhard - Olgemöller - Fricke,
Patentanwälte,
Postfach 10 09 62
80083 München (DE)

(54) **Modified lupin proteins for the preparation of water dispersible product forms of fat soluble compounds**

(57) The present invention relates to a process for the manufacture of a modified lupin protein, which comprises adjusting an aqueous solution or suspension of lupin proteins of native origin having a dry mass content of from 0,1 to 20 % to pH 3 to 9, adding 0,01 to 10 % by weight of a protease, in relation to the dry weight of the lupin protein, incubating the protein solution or suspension at a temperature of from 5 to 70 °C until a degree of hydrolysis of from 1 to 30 % is obtained, and inacti-

vating the protease, to modified lupin proteins obtainable by this process, to compositions containing said lupin proteins together with a fat-soluble active ingredient or colorant, and to the use of said composition for enrichment, fortification and/or coloration of food, beverages, animal feeds, cosmetics or drugs. Further, the invention relates to food, beverage, animal feeds, cosmetics or drugs containing said composition.

EP 1 405 572 A1

Description

[0001] The present invention relates to novel lupin proteins, a process for the manufacture thereof and compositions containing finely divided fat soluble actives and/or colorants in a matrix based on the novel lupin proteins and to a process for preparing these compositions. The present invention further relates to the use of the novel compositions of this invention for coloration and /or for enrichment or fortification of food, beverages, animal feeds, cosmetics or drugs. More particularly, the present invention relates to novel compositions comprising an enzymatically and/or physically treated lupin protein and a fat soluble active or colorant, to a process for preparing these compositions and the use of these compositions as additives for coloration and/or enrichment or fortification for food, beverages, animal feeds, cosmetics or drugs; and to food, beverages, animal feeds, cosmetics or drugs containing such compositions.

[0002] In one aspect, the present invention relates to a process for the manufacture of a modified lupin protein, which comprises enzymatically hydrolysing a lupin protein of native origin. More particularly, the invention relates to a process which comprises adjusting an aqueous solution or suspension of lupin proteins of native origin having a dry mass content of from 0,1 to 20 % to pH 3 to 9, adding 0,01 to 10 % by weight, in relation to the weight of the dry lupin protein, of a protease, incubating the protein solution or suspension at a temperature of from 5 to 70 °C until obtention of a degree of hydrolysis of from 1 bis 30 %, inactivating of the protease, and converting the hydrolysate into a solid form. If required the said protein hydrolysate may be physically fractionated by common methods such as centrifugation and/or ultrafiltration. In another aspect the present invention relates to a process which comprises adjusting an aqueous solution or suspension of lupin proteins of native origin having a dry mass content of from 0,1 to 20 % to pH 3 to 9, fractionating the solution or suspension by centrifugation and/or ultrafiltration and converting the protein fraction desired into a solid form. The solid product may either represent the water soluble or the insoluble fraction.

[0003] The term "fat soluble active" as used herein comprises a fat soluble vitamins or functionally related compounds which can be used for enrichment or fortification of food, beverages, animal feeds, cosmetics or drugs. Examples of such fat soluble vitamins or the vitamins of the groups A, D, E or K or derivatives thereof such as their acetates or their longer chain fatty acid esters. Examples for functionally related compounds are e.g. polyunsaturated fatty acids (PUFAs) or derivatives thereof, triglycerides rich in polyunsaturated fatty acids such as eicosapentaenoic acid (EPA), docosahexaenoic acid (DHA) or γ -linolenic acid (GLA), or coenzyme Q 10 (CoQ 10). Also included are fat soluble sun-filters, such as UV-A and UV-B filters used in sun care and cosmetic preparations.

[0004] The term "colorant" as used herein comprises a carotene or structurally related polyene compound which can be used as a colorant for food, beverages, animal feeds, cosmetics or drugs. Examples of such carotenoids are α - or β -carotene, 8'-apo- β -carotenal, 8'-apo- β -carotenoic acid esters such as the ethyl ester, canthaxanthin, astaxanthin, lycopene, lutein, zeaxanthin or crocetin, α - or β -zeacarotene or mixtures thereof. The preferred carotenoid is β -carotene. It is understood that the above named substances of the categories "fat soluble active" and "colorant" can also be used as mixtures within the novel compositions of the present invention.

[0005] The term "lupin protein of native origin" refers to proteins which can be isolated from seeds (including prepared or processed seeds) of any variety of the lupin plant (e.g., *L. albus*, *L. angustifolius* or varieties thereof), or from whole flour or defatted products such as shred or white flakes. Lupin protein concentrates and thermically or chemically pretreated lupin products may be used also for the purposes of the invention. In a preferred embodiment of the invention, the modified lupin proteins are obtained from solutions or suspension of native lupin protein which have a dry mass content of from about 5-15 % by weight. The solution or suspension is adjusted to pH 3-9, preferably pH 4-8, e.g., by the addition of 0.1-2 M acid or base such as hydrochloric acid and sodium hydroxide, respectively and preferably brought to a temperature of from 40-60 °C, especially 50 °C. The protease used may be a protease having endo-activity or exo-activity. Mixtures of endo- and exo-proteases may be used. Proteases for use in the process of the invention are available from the suppliers, Novozymes, Genencor, AB-Enzymes, Gist-Brocades etc. Preferred proteases are those of bacterial or fungal origin, e.g. from *Bacillus licheniformes* or *Aspergillus oryzae*. Enzyme preparations such as Alcalase or Flavourzyme as obtainable from the firm Novozymes may be used. The protease is added to provide a concentration of about 0.01 - 10 % by weight, preferably about 0.1 - 1 % by weight in relation to the weight of the lupin protein. The enzymatic hydrolysis is carried out until a degree of hydrolysis of about 1-30 %, preferably 1-10 % is reached. The degree of hydrolysis can be determined in a manner known to those skilled in the art, e.g. as described by Petersen, D., Nielsen, P.M., Cambmann C., Determination of the Degree of Hydrolysis (DH) based on OPA Reaction, ED-9512723, Novo Nordisk A/S, Dec. 1995; Frister, H., Meisel, H., Schlimme, E., OPA method modified by use of N,N-dimethyl-2-mercaptoethylammonium chloride as thiol component, Fresenius J. Anal. Chem. 330 (1988) 631. The expression "degree of hydrolysis" means the number of hydrolyzed peptide bonds in relation to the total sum of peptide bonds in the lupin protein originally present in the mixture.

[0006] The protease is then inactivated, suitably by heating of the incubation batch. In a preferred embodiment of the process of the invention, the enzyme is added in one single step. The enzymatic hydrolysis may also be carried out stepwise. For instance, the protease or a mixture of proteases is added to the incubation batch in an amount of e.g., 1 % whereupon, e.g., after 5-10 minutes (at a temperature of 50 °C) further protease or mixture of proteases which

may by the same or different from the first added protease or mixture of proteases is added, e.g. in an amount of 2 % whereupon the incubation batch is hydrolyzed at 50 ° for 10 minutes. Using this procedure, starting proteins having a degree of hydrolysis of approximately zero can be used.

[0007] Basically, all the above-mentioned parameters are variable. Thus, the temperature of the hydrolysis step may be, in general, 5-70 °C with temperature of 40-60 °C, especially 50 °C being preferred. The duration of hydrolysis may vary between 1 and 300 minutes and is preferably 5-10 minutes. The amount of enzyme added may vary between about 0,01-10 % by weight. When the hydrolysis is carried out in a two-step procedure, the amount of enzyme used in the first step is preferably 1 % by weight and in the second step 2 % by weight (based on the dry mass of protein).

[0008] The inactivation of the protease can be effected by heat denaturation, e.g., by heating to 80-85 °C for 5-10 minutes. Subsequently, the product may be submitted to fractionating, e.g., by centrifugation or ultrafiltration to separate and isolate the modified lupin protein.

[0009] The so-obtained solution or suspension of the modified lupin protein is then converted into a solid form, e.g. a dry powder, e.g., by spray-drying or freeze-drying.

[0010] In another aspect of the invention, the modified lupin protein may be obtained by adjusting an aqueous solution or suspension of lupin proteins having a dry mass content of from 0,1 to 20 %, preferably 5-15 % by weight to pH 3 to 9, preferably 4-6, fractionating the solutions or suspension, e.g., by procedures such as centrifugation or ultrafiltration and converting the fractionated hydrolysate into a solid form. Prior to drying, the modified lupin protein fractions may be submitted to heat treatment (Pasteurisation) e.g. by heating to 80-85 °C for 5-10 minutes to stabilize the product against microbial deterioration.

[0011] In still another aspect the present invention relates to modified lupin proteins which are obtainable by the process of the present invention, and to compositions comprising such modified lupin proteins and a fat-soluble active ingredient or colorant, and, optionally, adjuvants and/or excipients.

[0012] The modified lupin protein is a water-soluble or dispersible polypeptide exerting a protective role towards the inner phase of the dispersion i.e. the fat soluble active or colorant and, therefore, acts as a protective hydrocolloid. As a rule the hydrocolloid features surface activity and imparts viscosity, by virtue of which it stabilizes emulsions or suspensions in which it is present with the fat soluble active or colorant.

[0013] The compositions of the present invention are preferably solid, powdered compositions wherein the contained fat soluble active or colorant is finely dispersed in a matrix or carrier. The matrix or carrier contains or consists of modified lupin protein as a protective hydrocolloid and, optionally, one or more water soluble excipients and/or adjuvants conventionally used for the formulation of fat soluble actives or colorants.

[0014] In the compositions of the present invention, the amount of modified lupin protein is from about 0.5 to about 60.0 wt.-% and the amount of fat soluble active and/or colorant is from about 0.1 to about 80.0 wt.-% based on the total weight of the composition, all components totaling 100 %.

[0015] Suitably, the compositions of the present invention (further) contain one or more excipients and/or adjuvants selected from one or more each of a mono- di-, oligo- and polysaccharides, glycerol, triglycerides, water-soluble antioxidants and fat-soluble antioxidants. Solid compositions may in addition contain an anti-caking agent, such as silicic acid, and up to about 10 wt.%, as a rule about 2 to 5 wt.%, of water.

[0016] Examples of mono- and disaccharides which may be present in the compositions of the present invention are sucrose, invert sugar, glucose, fructose, lactose, maltose, saccharose and sugar alcohols, and examples of the oligo- and polysaccharides are starch and starch hydrolysates, e.g. dextrans and maltodextrans, especially those having the range of 5 - 65 dextrose equivalents (DE), and glucose syrup, especially such having the range of 20 - 95 DE. The term "dextrose equivalent" (DE) denotes the degree of hydrolysatation and is a measure of the amount of reducing sugar calculated as D-glucose based on dry weight; the scale is based on native starch having a DE close to 0 and glucose having a DE of 100.

[0017] The triglyceride is suitably a vegetable oil or fat, preferably corn oil, sunflower oil, soybean oil, safflower oil, rape seed oil, peanut oil, palm oil, palm kernel oil, cotton seed oil or coconut oil.

[0018] The organic solvent may be for example methylene chloride, chloroform, ethyl acetate, dimethyl ether, acetone, ethanol or isopropanol.

[0019] The water-soluble antioxidant may be for example ascorbic acid or a salt thereof, preferably sodium ascorbate. The fat-soluble antioxidant may be for example a tocopherol, e.g. dl- α -tocopherol (i.e. synthetic tocopherol), d- α -tocopherol (i.e. natural tocopherol), β - or γ -tocopherol, or a mixture of two or more of these; butylated hydroxytoluene (BHT); butylated hydroxyanisole (BHA); ethoxyquin, propyl gallate; tert. butyl hydroxyquinoline; or an ascorbic acid ester of a fatty acid, preferably ascorbyl palmitate or stearate. Depending on the pH of the aqueous matrix solution the ascorbic acid ester of a fatty acid, particularly ascorbyl palmitate or stearate, may alternatively be added to the water phase.

[0020] The compositions of the present invention may be solid compositions, i.e. stable, water-soluble or -dispersible powders, or they may be liquid compositions, i.e. aqueous colloidal solutions or oil-in-water dispersions of the aforementioned powders. The stabilized oil-in-water dispersions, which may be oil-in-water emulsions or may feature a

mixture of suspended, i.e. solid, particles and emulsified, i.e. liquid, droplets, may be prepared by the methods already described above.

[0021] Typically, a powder composition according to the present invention comprises

- 5 - about 0.5 to about 60 wt.-%, preferably about 5 to about 50 wt.-% of a modified lupin protein;
- about 0.1 to about 80 wt.-% preferably about 0.5 to about 60 wt.-% of a fat soluble active and/or a colorant;
- 0 to about 70 wt.-% preferably about 0 to about 40 wt.-% of a mono- or disaccharide;
- 0 to about 70 wt.-% preferably about 0 to about 40 wt.-% of a starch hydrolysate;
- 0 to about 20 wt.-% preferably about 0 to about 10 wt.-% of glycerol;
- 10 - 0 to about 50 wt.-% preferably about 0 to about 30 wt.-% of a triglyceride;
- 0 to about 5 % preferably about 0 to about 2 wt.-% of one or more water-soluble antioxidant(s);
- 0 to about 5 % preferably about 0 to about 2 wt.-% of one or more fat-soluble antioxidant(s);
- 0 to about 50 wt.-% preferably about 0 to about 35 wt.-% of a starch;
- 0 to about 5 wt.-% preferably about 1 wt.-% of silicic acid; and
- 15 - 0 to about 10 wt.-% preferably about 1 to about 5 wt.-% of water;

[0022] the percentages of all ingredients totaling 100.

[0023] The novel compositions of this invention can be used for coloration and /or for enrichment or fortification for food, beverages, animal feeds, cosmetics or drugs.

20 [0024] By the present invention there are preferably provided compositions comprising β -carotene as a colouring agent. These compositions, when dissolved, dispersed or diluted in/with water to a final β -carotene concentration of 10 ppm are typically characterized by ultraviolet/visible-spectroscopy using deionized water as reference. At a sample thickness of 1 cm the dispersions show an extinction of at least 0.6 (preferably above 1.0) absorbance units at the wavelength of maximum optical density in the range of 400 to 600 nm. This is equivalent to a formal extinction coefficient

25 of β -carotene in aqueous dispersion $E(1\%, 1 \text{ cm})$ of 600 (preferably >1000).

[0025] The manufacture of the compositions of the present invention can be carried out in a manner known per se for the preparation matrix-based compositions of fat soluble actives or colorant compositions for coloration and /or for enrichment or fortification for food, beverages, animal feeds, cosmetics or drugs, e.g. as disclosed in European Patent Publications Nos. 0 285 682, 0 347 751, 0 966 889, 1066 761 and 1 106 174, in the PCT Publication No. WO 98/15195

30 and in Swiss Patent Application No. 1238/92, the contents of which are incorporated herein by reference.

[0026] Thus, the compositions of the present invention can be prepared by a process which comprises homogenizing, in an aqueous or colloidal solution of the modified lupin protein and optionally one or more water-soluble excipients and/or adjuvants, a fat soluble active or colorant or a solution or dispersion of the fat soluble active or colorant and optionally one or more fat-soluble excipients and/or adjuvants, and converting the so-obtained dispersion into a solid

35 composition.
[0027] Typically, modified lupin protein and, optional water-soluble excipients and adjuvants are dissolved or dispersed in water. The fat soluble active and/or colorant and, optional fat-soluble excipients and adjuvants are either used as such or dissolved or suspended in a triglyceride and/or an organic solvent. The fat soluble active and/or colorant or the solution or dispersion thereof, respectively, is then added to the aqueous protein solution with stirring and the mixture homogenized using conventional technology. The so-obtained oil-in-water dispersion can, after removal of the organic solvent (if present), be converted into a solid composition, e.g. a dry powder.

40 [0028] A further object of the invention is a process for the preparation of the compositions based on a modified lupin protein as described in more detail above which process comprises homogenizing, in an aqueous solution or colloidal solution of the modified lupin protein and optional water-soluble excipients and adjuvants, a solution or dispersion of the fat soluble active and/or colorant and optional fat-soluble adjuvants and, if required, converting the dispersion obtained into a powder.

45 [0029] For the homogenization conventional technologies, such as high pressure homogenization, high shear emulsification (rotor-stator systems), micronization or wet milling, can be applied. Other techniques used for the preparation of fat soluble active or colorant compositions for coloration and /or for enrichment or fortification for food, beverages, animal feeds, cosmetics or drugs are disclosed in European Patent Publications Nos. 937412 and 1008380 and in US Patent Specification No. 6,093,348, the contents of which are incorporated herein by reference.

50 [0030] The so-obtained dispersion, which is an oil-in-water dispersion, can be converted into a solid composition, e.g. a dry powder, using any conventional technology such as spray drying, spray drying in combination with fluidized bed granulation (the latter technique commonly known as fluidized spray drying or FSD); or by a powder-catch technique whereby sprayed emulsion droplets are caught in a bed of an absorbent, such as starch, and subsequently dried.

55 [0031] Beverages wherein the product forms of the present invention can be used as a colorant or a functional ingredient can be carbonated beverages e.g., flavored seltzer waters, soft drinks or mineral drinks, as well as non-carbonated beverages e.g. flavored waters, fruit juices, fruit punches and concentrated forms of these beverages. They

may be based on natural fruit or vegetable juices or on artificial flavors. Also included are alcoholic beverages and instant beverage powders. Besides, sugar containing beverages diet beverages with non-caloric and artificial sweeteners are also included.

[0032] Further, dairy products, obtained from natural sources or synthetic, are within the scope of the food products wherein the product forms of the present invention can be used as a colorant or as a functional ingredient. Typical examples of such products are milk drinks, ice cream, cheese, yoghurt and the like. Milk replacing products such as soy milk drinks and tofu products are also comprised within this range of application.

[0033] Also included are sweets which contain the product forms of the present invention as a colorant or as a functional ingredient, such as confectionery products, candies, gums, desserts, e.g. ice cream, jellies, puddings, instant pudding powders and the like.

[0034] Also included are cereals, snacks, cookies, pasta, soups and sauces, mayonnaise, salad dressings and the like which contain the product forms of the present invention as a colorant or a functional ingredient. Furthermore, fruit preparations used for dairy and cereals are also included.

[0035] The final concentration of the active ingredient or the colorant which is added via the product forms of the present invention to the food products may be from about 0.1 to about 500 ppm, particularly from about 1 to about 50 ppm based on the total weight of the food composition and depending on the particular food product to be colored or fortified and the intended grade of coloration or fortification.

[0036] The food compositions of this invention are preferably obtained by adding to a food product the active ingredient or the colorant in the form of a composition of this invention. For coloration or fortification of a food or a pharmaceutical product a composition of this invention can be used according to methods per se known for the application of water dispersible solid product forms.

[0037] In general the composition may be added either as an aqueous stock solution, a dry powder mix or a preblend with other suitable food ingredients according to the specific application. Mixing can be done e.g. using a dry powder blender, a low shear mixer, a high pressure homogenizer or a high shear mixer depending on the formulation of the final application. As will be readily apparent such technicalities are within the skill of the expert.

[0038] Pharmaceutical compositions such as tablets or capsules wherein the compositions are used as a colorant are also within the scope of the present invention. The coloration of tablets can be accomplished by adding the product forms in form of a liquid or solid colorant composition separately to the tablet coating mixture or by adding a colorant composition to one of the components of the tablet coating mixture. Colored hard or soft shell capsules can be prepared by incorporating a colorant composition in the aqueous solution of the capsule mass.

[0039] Pharmaceutical compositions such as tablets such as chewable tablets, effervescent tablets or filmcoated tablets or capsules such as hard shell capsules wherein the compositions are used as an active ingredient are also within the scope of the present invention. The product forms are typically added as powders to the tableting mixture or filled into the capsules in a manner per se known for the production of capsules.

[0040] Animal feed products such as premixes, compound feeds, milk replacers, liquid diets or feed preparations wherein the compositions are either used as a colorant for pigmentation e.g. for egg yolks, table poultry, broilers or aquatic animals or as an active ingredient are also within the scope of the present invention.

[0041] Cosmetics, toiletries and derma products i.e. skin and hair care products such as creams, lotions, baths, lipsticks, shampoos, conditioners, sprays or gels wherein the compositions are used as a colorant or as an active ingredient are also within the scope of the present invention.

[0042] For the production of liquid and solid product forms such as oil-in-water suspensions, oil-in-water emulsions or powders the hydrocolloids used therein act as multifunctional ingredients. For this purpose these hydrocolloids should show a spectrum of functional properties such as excellent dispersion stabilizing properties, high emulsifying capacity and efficiency (the latter being relevant with respect to the particle size distribution of the inner phase of the dispersion), high solubility and well adapted viscosity in aqueous solution, thermal stability, low susceptibility to pH, ion strength and charge, superior film-forming as well as acceptable gas and water vapour barrier properties in solid product forms.

[0043] Native lupin proteins are readily commercially available in the form of flours or protein concentrates such as a product Vitaprot Type LP 60 of J. Rettenmaier & Sons.

[0044] The steps used for modification of the lupin protein isolate described above lead to overall improved functional properties of the protein such as better emulsifying properties, generally higher solubility in aqueous solution e.g. lower susceptibility to pH and counter ions as well as better cold water solubility, better temperature resistance, and better film-forming properties. Furthermore the viscosity in aqueous solution is decreased significantly and is now in an acceptable range for the purpose of the current invention.

[0045] The following Examples illustrate the invention further

Example 1

[0046] An aqueous suspension of native lupin protein is prepared by aqueous extraction of defatted lupin seed at pH 4-5 and subsequent protein extraction at pH 6-8. The aqueous suspension of native lupin protein has a dry matter content of 4 % by weight the dry matter having a protein content of > 90 %. The pH of the suspension is then adjusted to pH 7 and to a temperature of 50°C. Thereafter, 0,5 % by weight of Alcalase 2.4 L (Novozymes) is added. During the enzymatic hydrolysis the pH is maintained constant by the addition of 1 M NaOH solution and the temperature is maintained at 50 °C. After 5-10 minutes the protease is inactivated by heating of the suspension to 80 °C for 10 minutes. Then the insoluble fraction is separated and the solution is spray-dried at a product temperature of 70-80 °C.

Example 2

[0047] An aqueous suspension of native lupin protein is prepared by aqueous extraction of defatted lupin seed at pH 4-5 (the soluble fraction representing the pre-extract) and subsequent protein extraction at pH 6-8. The aqueous suspension of native lupin protein has a dry matter content of 4 % by weight the dry matter having a protein content of > 90 %. The suspension is adjusted to pH 4-5 and the insoluble fraction is separated by centrifugation. The supernatant is combined with the acidic pre-extract and the combined solution is purified by ultrafiltration. Thereby, the soluble protein components of the acidic pre-extract and the soluble proteins of the acidified protein extract in the retentate are obtained. The retentate of the ultrafiltration is spray-dried at a product temperature of 70-80 °C.

Example 3

[0048] A composition comprising modified lupin protein and vitamin A is prepared as follows:

a) Preparation of (aqueous) solution A:

[0049] A dry premix of 34.1 g of modified lupin protein L143T (prepared from L. albus Typ Top according to Example 1 and referred to hereinafter as L143T) and 51.2 g of dried glucose syrup Glucidex IT 47 (Roquette Frères, F-Lestrem) was prepared and subsequently mixed with 120 ml of deionized water in a 1 l reaction vessel at room temperature. The mixture was heated to 45°C and stirred for about 15 minutes at 45°C. The pH was around 6.1.

b) Preparation of (oil-based) solution B:

[0050] 15.3 g of crystalline vitamin A acetate, 2.7 g of peanut oil and 1.3 g of dl- α -tocopherol were introduced into a 100 ml Erlenmayer flask. A magnetic stirrer bar was added to the mixture. While stirring the mixture was heated to 65°C within 10 minutes.

c) Preparation of emulsion from solutions A and B:

[0051] Solution A was brought to 50°C and while vigorously stirring solution B was added to solution A within 3 minutes and the dispersion was vigorously stirred for another 20 minutes at 50°C. The emulsion was heated to 60°C and stirred for 5 hours in order to investigate its long term physical stability. After the holding time the mean particle size of the inner phase of the emulsion was 388 nm as measured by photon correlation spectroscopy (Coulter N4 Plus).

d) Preparation of solid composition from the emulsion:

[0052] The emulsion was sprayed into a pre-cooled fluidized bed of corn starch. Excess corn starch was removed by sieving and the powder obtained was dried in an air stream at room temperature for about 2 hours. The powder particle fraction (126.5 g) in the range of 0.16-0.50 mm was collected by sieving and characterized with respect to vitamin A content, surface oil contamination, corn starch content and residual humidity. The powder obtained had a vitamin A content of 263'000 IU/g as measured by HPLC analysis. The surface oil contamination was 0.2 % of the vitamin A content. The powder had a corn starch content of 24.4 % and a residual water content of 6.6 %.

Example 4

[0053] A composition comprising modified lupin protein and vitamin E is prepared as follows:

a) Preparation of (oil-based) solution A:

[0054] 26.7 g of dl- α -tocopheryl acetate were introduced into a 100 ml Erlenmeyer flask. A magnetic stirrer bar was added and the dl- α -tocopherylacetate was heated to 75°C.

b) Preparation of (aqueous) solution B:

[0055] A dry premix of 40.0 g of modified lupin protein L143T was prepared and subsequently mixed with 135 ml of deionized water in a 1 l reaction vessel at room temperature. The mixture was heated to 50°C and stirred for about 15 minutes at 50°C. The pH was around 6.1.

c) Preparation of emulsion from solutions A and B:

[0056] Solution B was brought to 55°C and while vigorously stirring solution A was added to solution B and the emulsion was vigorously stirred for another 30 minutes at 55°C. The emulsion was heated to 60°C and stirred for 5 hours in order to investigate its long term physical stability. After the holding time the mean particle size of the inner phase of the emulsion was 521 nm as measured by photon correlation spectroscopy (Coulter N4 Plus).

d) Preparation of solid composition from the emulsion:

[0057] The emulsion was diluted with deionized water to a final water content of 65 % and then spray dried with a Mobile Minor (Niro/VS). The inlet temperature of the drying air was 220°C the outlet temperature was 110°C. The powder was collected and about 1% of silicon dioxide was added. The tocopheryl acetate content of the powder was 25.6 % as measured by HPLC.

Example 5

[0058] A composition comprising modified lupin protein and β -carotene is prepared as follows:

a) Preparation of (oil-based) solution A:

[0059] 8.8 g of corn oil and 1.6 g of dl- α -tocopherol were introduced into a 500 ml Erlenmeyer flask. 18.4 g of crystalline β -carotene were dispersed in 206 ml of trichloro methane and the dispersion obtained was added to the corn oil and the tocopherol. A magnetic stirrer bar was added to the mixture and a reflux condenser was fitted to the Erlenmeyer flask. By gently stirring and, at the same time, heating the mixture to about 65°C a solution was obtained.

b) Preparation of (aqueous) solution B:

[0060] A dry premix of 48.0 g of lupin protein L140T-2 (prepared from *L. angustifolius* according to Example 1 and referred to hereinafter as L140T-2) and 40.8 g of sucrose and 2.4 g of ascorbyl palmitate was prepared and subsequently mixed with 140 ml of deionized water in a 1 l reaction vessel at room temperature. While stirring the mixture was heated to 35°C and by adding 14 ml of a 1 N sodium hydroxide solution the pH of the mixture was brought to 7.9. Subsequently, the mixture was vigorously stirred for a short time in order to achieve a homogeneous solution.

c) Preparation of emulsion from solutions A and B:

[0061] While vigorously stirring, solution A was added to solution B at 35°C and the dispersion was vigorously stirred for another 30 minutes. The reaction vessel was flushed with nitrogen and the dispersion was heated to 55°C. The stirred dispersion was kept at 55°C for 60 minutes. Residual trichloro methane was removed at 50-55°C using a rotary evaporator. After removing entrapped air bubbles by centrifugation the emulsion was gently stirred at 50-55°C for some minutes and then characterized with respect to particle size of the inner phase. The mean particle size of the inner phase of the emulsion was 148 nm, as measured by photon correlation spectroscopy (Coulter N4 Plus).

d) Preparation of solid composition from the emulsion:

[0062] The emulsion was sprayed into a pre-cooled fluidized bed of corn starch. Excess corn starch was removed by sieving and the powder obtained was dried in an air stream at room temperature for about 2 hours. The powder particle fraction (approx. 138 g) in the range of 0.16-0.63 mm was collected by sieving and characterized with respect

to carotenoid content, colour intensity and colour hue in aqueous dispersion, content of corn starch and residual humidity.

[0063] The obtained powder had a β -carotene content of 12.8 % (11.5 %), as measured by UV/VIS-spectroscopy (HPLC), a content of 25 % corn starch and a residual water content of 5.9 %. It was dispersed in deionized water and the extinction was measured in a 1 cm quartz precision cell against deionized water. For a 10 ppm dispersion of β -carotene an extinction of 1.130 at a wavelength of 481 nm was calculated [$E(1\%, 1\text{ cm}) = 1130$]. The colour values $L^* = 81.9$, $a^* = 22.6$ and $b^* = 31.6$ were measured according to the CIE-system for a 5 ppm dispersion of β -carotene with a Hunterlab Ultrascan Spectrocolorimeter (1 cm, TTRAN). Based upon the values of a^* and b^* a colour hue angle $h^* = 54.4^\circ$ at a saturation (chroma value) $C^* = 38.8$ can be calculated.

Example 6

[0064] A composition comprising modified lupin protein and β -carotene is prepared as follows:

[0065] The procedure according to Example 5 was carried out using modified lupin protein L143T.

[0066] The obtained powder had a β -carotene content of 14.2 % (12.4 %), as measured by UV/VIS-spectroscopy (HPLC), a content of 20 % corn starch and a residual water content of 5.5 %. It was dispersed in deionized water and the extinction was measured in a 1 cm quartz precision cell against deionized water. For a 10 ppm dispersion of β -carotene an extinction of 0.892 at a wavelength of 478 nm was calculated [$E(1\%, 1\text{ cm}) = 892$]. The colour values $L^* = 84.1$, $a^* = 16.8$ and $b^* = 27.4$ were measured according to the CIE-system for a 5 ppm dispersion of β -carotene with a Hunterlab Ultrascan Spectrocolorimeter (1 cm, TTRAN). Based upon the values of a^* and b^* a colour hue angle $h^* = 58.5^\circ$ at a saturation (chroma value) $C^* = 32.1$ can be calculated.

Example 7

[0067] A composition comprising modified lupin protein and β -carotene is prepared as follows:

a) Preparation of (aqueous) solution A:

[0068] 60 g of modified lupin protein L143T were dissolved in 400 ml of deionized water at 50°C . A dry premix consisting of 64.5 g of sucrose, 121.8 g of a maltodextrin (DE 20-23) and 3.0 g sodium ascorbate was prepared and added to the aqueous protein solution at 50°C (= solution A).

b) Preparation of (oil-based) solution B:

[0069] 46.5 g of a triglyceride (Durkex 500: partly hydrogenated soybean oil) and 0.3 g of dl- α -tocopherol were mixed and heated to 170°C . Subsequently, 3.75 g of β -carotene was suspended in the mixture of triglyceride and tocopherol. By stirring for about 10 minutes a clear solution of β -carotene was obtained (= solution B).

c) Preparation of emulsion from solutions A and B:

[0070] A crude emulsion was prepared by adding solution B to solution A while gently stirring. A fine emulsion was obtained by a four passage high pressure homogenizing treatment of the preemulsion at a pressure of 50/400 bar (LAB 1000 of APV).

d) Preparation of solid composition from the emulsion:

[0071] The emulsion was spray dried in a laboratory spray dryer (Mobile Minor of Niro) at an inlet temperature of 190°C and an outlet temperature of 83°C . A fine powder was obtained with a residual water content of 1.8 %. The β -carotene content of the powder was 1.3 % as measured by UV/VIS-spectroscopy and HPLC. The powder was dispersed in deionized water and the extinction was measured in a 1 cm quartz precision cell against water. For a 10 ppm dispersion of β -carotene an extinction of 1.895 at a wavelength of 463 nm was calculated [$E(1\%, 1\text{ cm}) = 1895$]. The colour values $L^* = 85.0$, $a^* = 9.0$ and $b^* = 63.9$ were measured according to the CIE-system for a 5 ppm dispersion of β -carotene with a Hunterlab Ultrascan Spectrocolorimeter (1 cm, TTRAN). Based upon the values of a^* and b^* a colour hue angle $h^* = 82.0^\circ$ at a saturation (chroma value) $C^* = 64.5$ can be calculated.

Example 8

[0072] A Soft Drink formulation with 6 ppm β -carotene can be prepared as follows :

Formula	
Ingredients	[g]
Sugar syrup 64°Brix	156.2
Na-benzoate	0.2
Ascorbic acid	0.2
Citric acid (as aqueous solution 50% w/w)	5.0
Flavor *)	0.5
β -carotene 10% as a 1 % stock solution **)	6.0
Water	to 1000.0

*) Apricot Flavour No. 78511-76 of Givaudan

**) i.e., a 1 % aqueous solution (dispersion) of formulations as obtained e.g. in Example 3 or 4, based on a β -carotene content of 10 % in said formulation

Procedure:

[0073] Dissolve Na-benzoate in part of the water, add sugar syrup, ascorbic acid, citric acid and flavor and finally the β -carotene stocksolution. Dilute the bottling syrup to one liter of beverage.

[0074] The β -carotene provides a vivid orange color of a high clarity and with a high coloring strength. Longtime stability tests of three months at ambient conditions showed a good physical stability of β -carotene in beverages and a good color stability in terms of color values $L^*C^*h^*$.

Claims

1. Process for the manufacture of a modified lupin protein, which comprises adjusting an aqueous solution or suspension of lupin proteins of native origin having a dry mass content of from 0,1 to 20 % to pH 3 to 9, adding 0,01 to 10 % by weight of a protease, in relation to the dry weight of the lupin protein, incubating the protein solution or suspension at a temperature of from 5 to 70 °C until a degree of hydrolysis of from 1 to 30 % is obtained, and inactivating the protease.
2. A process according to claim 1, wherein the pH of the solution or suspension is adjusted to pH 4 to 8, the amount of protease added is 0,1 to 1 %, and the enzymatic hydrolysis is effected at 20 to 55 °C.
3. A process according to claims 1 or 2, wherein the enzymatic hydrolysis is effected to a degree of hydrolysis of from 1 to 10 %.
4. A process according to any one of claims 1 to 3, wherein a protease having endo-activity is used.
5. A process according to any one of claims 1 to 3, wherein a protease having exo-activity is used.
6. A process according to any one of claims 1 to 3, wherein a protease from bacteria or fungi is used.
7. A process according to any of the preceding claims, wherein the modified lupin protein, after the protease has been inactivated, is brought into a solid form by centrifugation or ultrafiltration, optionally after a fractionation step.
8. Process for the manufacture of a modified lupin protein, which comprises adjusting an aqueous solution or suspension of lupin proteins of native origin having a dry mass content of from 0,1 to 20 % to pH 3 to 9, fractionating the solutions or suspension by centrifugation or ultrafiltration, and converting the protein solution into a solid form.
9. Modified lupin proteins obtainable according to the process of any one of claim 1-8.
10. Composition, comprising a modified lupin protein according to any one of claims 1-8, and a fat-soluble active ingredient or colorant, and, optionally, adjuvants and/or excipients.

11. A composition as in claim 10 wherein the fat soluble active ingredient or colorant is a carotenoid, a fat soluble vitamin, a triglyceride, an oil soluble UV-A or UV-B filter or a mixture thereof.
12. A composition as in claim 11 wherein the carotenoid is α - or β -carotene, 8'-apo- β -carotenal, 8'-apo- β -carotenoic acid ethyl ester, canthaxanthin, astaxanthin, lycopene, lutein, zeaxanthin or crocetin or mixtures thereof.
13. A composition as in claim 11 wherein the fat soluble vitamin is Vitamin A, D, E or K.
14. A composition as in any one of claims 9-13 wherein at least one of a mono- di-, oligo- or polysaccharide, a triglyceride, a water-soluble anti-oxidant, a fat-soluble anti-oxidant, silicic acid and water is additionally present.
15. A composition as in claim 14 wherein the mono- or disaccharide is saccharose, invert sugar, glucose, fructose, lactose or maltose and the composition optionally contains glycerol in addition.
16. A composition as in claim 14 wherein the polysaccharide is a starch or a starch hydrolysate and /or the triglyceride is a vegetable oil or fat.
17. A composition as in claim 16 wherein the starch hydrolysate is a dextrin or a maltodextrin (in the range of 5-65 dextrose equivalents) or a glucose syrup (in the range of 20-95 dextrose equivalents).
18. A composition as in any one of claims 10-17 wherein the amount of modified lupin protein is from about 0.5 to about 60.0 wt.-% and the amount of fat soluble active ingredient or colorant is from about 0.1 to about 80.0 wt.-%.
19. A process for the preparation of a composition as claimed in any one of claims 10-18 which comprises homogenizing, in an aqueous solution or colloidal solution of the modified lupin protein and optional water-soluble excipients and adjuvants, a solution or dispersion of the fat soluble active and/or colorant and optional fat-soluble adjuvants and, if required, converting the dispersion obtained into a powder.
20. Use of a composition as claimed in any one of claims 10-18 for enrichment, fortification and/or coloration for food, beverages, animal feeds, cosmetics or drugs.
21. Food, beverages, animal feeds, cosmetics or drugs containing a composition as claimed in any one of claims 10-18.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 02 02 2479

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
A	WO 99 51106 A (BUTTIMER EILEEN TERESA; FITCHETT COLIN STANLEY (GB); HOWARD JULIE) 14 October 1999 (1999-10-14)	1-7	A23L1/303 A23J3/34 A23L1/275 A23L1/30
X	* page 9, line 4-23; claims 1-51 *	8-10, 21	
Y	* examples 1-5 *	11-20	
A	EP 0 859 553 A (FINANZCONSUL AG; FRAUNHOFER GES FORSCHUNG (DE)) 26 August 1998 (1998-08-26)	1-7, 10, 21	
X	* column 7, paragraph 1 *	8, 9	
	* column 4, line 32-36 *		
	* column 4, line 47-58 *		
	* column 11, line 9 - column 12, line 4; claim 5 *		
Y		11-20	
A	DE 100 64 434 A (JAEGLLE WOLFGANG) 11 July 2002 (2002-07-11)	1-7, 10, 21	
X	* the whole document *	8, 9	
Y		11-20	
A	US 6 335 044 B1 (LUCK THOMAS ET AL) 1 January 2002 (2002-01-01)	1-21	TECHNICAL FIELDS SEARCHED (Int.Cl.7) A23J A23L
	* the whole document *		
A	ALAMANOU S ET AL: "EMULSIFYING PROPERTIES OF LUPIN SEED PROTEINS (LUPINUS ALBUS, SSP. GRAECUS)" DEVELOPMENTS IN FOOD SCIENCE, ELSEVIER, AMSTERDAM, NL, vol. 37B, 6 July 1995 (1995-07-06), pages 2129-2138, XP002068534 ISSN: 0167-4501	1-21	
	* the whole document *		
A	EP 1 219 292 A (BASF AG) 3 July 2002 (2002-07-03)	1-21	
	* the whole document *		
-/--			
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 5 August 2003	Examiner De Jong, E
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document</p> <p>T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons</p> <p>& : member of the same patent family, corresponding document</p>			

EPO FORM 1503 D3 82 (10/04C01)



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 02 02 2479

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (InCL.7)
A	EP 1 106 174 A (HOFFMANN LA ROCHE) 13 June 2001 (2001-06-13) * the whole document *	1-21	
A	EP 0 937 412 A (HOFFMANN LA ROCHE) 25 August 1999 (1999-08-25) * claims 1-11 *	1-21	
A	EP 0 347 751 A (HOFFMANN LA ROCHE) 27 December 1989 (1989-12-27) * claims 1-10 *	1-21	
A	US 4 522 743 A (DITTER WALTER ET AL) 11 June 1985 (1985-06-11) * claims 1-5; examples 1-17 *	1-21	
A	WO 02 19836 A (CORP CRYOLAB ;MCNEIL MARCEL C (CA)) 14 March 2002 (2002-03-14) * page 8, line 26-34; claims 1-11 *	1-21	
A	RAYMUNDO A ET AL: "OPTIMIZATION OF THE COMPOSITION OF LOW-FAT-OIL-IN-WATER EMULSIONS STABILIZED BY WHITE LUPIN PROTEIN" JOURNAL OF THE AMERICAN OIL CHEMISTS' SOCIETY, AMERICAN OIL CHEMISTS' SOCIETY, CHAMPAIGN, US, vol. 79, no. 8, August 2002 (2002-08), pages 783-790, XP001124618 ISSN: 0003-021X * page 783, column 2, paragraph 3; table 1 *	1-21	<div>TECHNICAL FIELDS SEARCHED (InCL.7)</div>
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 5 August 2003	Examiner De Jong, E
CATEGORY OF CITED DOCUMENTS X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document		T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons & : member of the same patent family, corresponding document	

EPO FORM 1503 03.02 (P04C01)



European Patent
Office

Application Number
EP 02 02 2479

CLAIMS INCURRING FEES

The present European patent application comprised at the time of filing more than ten claims.

☐ Only part of the claims have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims and for those claims for which claims fees have been paid, namely claim(s):

☐ No claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims.

LACK OF UNITY OF INVENTION

The Search Division considers that the present European patent application does not comply with the requirements of unity of invention and relates to several inventions or groups of inventions, namely:

see sheet B

☒ All further search fees have been paid within the fixed time limit. The present European search report has been drawn up for all claims.

☐ As all searchable claims could be searched without effort justifying an additional fee, the Search Division did not invite payment of any additional fee.

☐ Only part of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the inventions in respect of which search fees have been paid, namely claims:

☐ None of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the invention first mentioned in the claims, namely claims:



European Patent
Office

LACK OF UNITY OF INVENTION
SHEET B

Application Number
EP 02 02 2479

The Search Division considers that the present European patent application does not comply with the requirements of unity of invention and relates to several inventions or groups of inventions, namely:

1. Claims: 1-7, 9-21 partly

- Process for the manufacture of a modified lupin protein, comprising hydrolysis by a protease until a degree of hydrolysis of 1-30%.
- Modified lupin protein (partially hydrolysed).
- Composition comprising a modified lupin protein (partially hydrolysed) and a fat-soluble active ingredient or colorant.
- Process for preparing the composition.
- Use of the composition.
- Food, beverage, animal feed, cosmetic or drug containing the composition.

2. Claims: 8, 9-21 (partly)

- Process for the manufacture of a a modified lupin protein, comprising fractionation by centrifugation or ultrafiltration and conversion into a solid form.
- Modified lupin protein (fractionated and dried).
- Composition comprising a modified lupin protein (fractionated and dried) and a fat-soluble active ingredient or colorant.
- Process for preparing the composition.
- Use of the composition.
- Food, beverage, animal feed, cosmetic or drug containing the composition.

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 02 02 2479

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

05-08-2003

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
WO 9951106	A	14-10-1999	AU	3218999 A	25-10-1999
			CN	1295440 T	16-05-2001
			EP	1065940 A1	10-01-2001
			WO	9951106 A1	14-10-1999
EP 0859553	A	26-08-1998	AT	192283 T	15-05-2000
			AU	717831 B2	06-04-2000
			AU	7619596 A	28-04-1997
			DE	19680849 D2	12-05-1999
			DE	59605147 D1	08-06-2000
			EP	0859553 A1	26-08-1998
			GR	3034113 T3	30-11-2000
			WO	9712524 A2	10-04-1997
			DE	19640992 A1	10-04-1997
			ES	2148813 T3	16-10-2000
			PT	859553 T	31-10-2000
DE 10064434	A	11-07-2002	DE	10064434 A1	11-07-2002
US 6335044	B1	01-01-2002	DE	19813207 C1	24-06-1999
			AT	218281 T	15-06-2002
			AU	738948 B2	27-09-2001
			AU	1332299 A	27-04-1999
			CA	2305287 A1	15-04-1999
			WO	9917619 A1	15-04-1999
			DE	59804355 D1	11-07-2002
			DK	1024706 T3	08-07-2002
			EP	1024706 A1	09-08-2000
			ES	2178284 T3	16-12-2002
			HR	20000197 A1	31-08-2000
			HU	0004826 A2	28-06-2001
			PL	339737 A1	02-01-2001
			PT	1024706 T	31-10-2002
			SK	4772000 A3	12-02-2001
			DE	19912037 A1	20-06-2002
EP 1219292	A	03-07-2002	DE	10064387 A1	27-06-2002
			CN	1362406 A	07-08-2002
			EP	1219292 A1	03-07-2002
			JP	2002262824 A	17-09-2002
			US	2002165285 A1	07-11-2002
EP 1106174	A	13-06-2001	EP	1106174 A1	13-06-2001
			AU	7201500 A	14-06-2001
			BR	0005801 A	27-11-2001
			CA	2328025 A1	09-06-2001

EPO FORM P4089

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

ANNEX TO THE EUROPEAN SEARCH REPORT ON EUROPEAN PATENT APPLICATION NO.

EP 02 02 2479

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

05-08-2003

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 1106174	A		CN 1300562 A	27-06-2001
			JP 2001172172 A	26-06-2001
			NO 20006182 A	11-06-2001
			US 2001009679 A1	26-07-2001
EP 0937412	A	25-08-1999	EP 0937412 A1	25-08-1999
			AU 743535 B2	31-01-2002
			AU 1836299 A	09-09-1999
			BR 9900776 A	28-03-2000
			CN 1231843 A	20-10-1999
			JP 2000186224 A	04-07-2000
			NO 990852 A	24-08-1999
			US 2001008644 A1	19-07-2001
EP 0347751	A	27-12-1989	AT 87472 T	15-04-1993
			AU 623908 B2	28-05-1992
			AU 3657389 A	04-01-1990
			CA 1340242 C	15-12-1998
			DE 58903922 D1	06-05-1993
			DK 310989 A	24-12-1989
			EP 0347751 A1	27-12-1989
			ES 2039759 T3	01-10-1993
			FI 893022 A	24-12-1989
			FI 981654 A	23-07-1998
			GR 3031153 T3	31-12-1999
			HK 142595 A	15-09-1995
			HU 52341 A2	28-07-1990
			IE 62542 B1	08-02-1995
			IL 90651 A	21-02-1993
			JP 2051594 A	21-02-1990
			JP 2542697 B2	09-10-1996
			KR 135759 B1	23-04-1998
			MC 2034 A	30-05-1990
			NO 892588 A ,B,	27-12-1989
			NZ 229599 A	28-10-1992
			PH 27096 A	26-02-1993
			PT 90948 A ,B	29-12-1989
			US 5478569 A	26-12-1995
			ZA 8904615 A	28-02-1990
US 4522743	A	11-06-1985	DE 3119383 A1	02-12-1982
			AT 16814 T	15-12-1985
			AU 553623 B2	24-07-1986
			AU 8372582 A	18-11-1982
			CA 1202809 A1	08-04-1986
			DE 3267761 D1	16-01-1986

EPO FORM P459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 02 02 2479

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

05-08-2003

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 4522743	A		DK 216082 A ,B,	16-11-1982
			DK 334188 A ,B,	17-06-1988
			EP 0065193 A2	24-11-1982
			IL 65707 A	31-03-1985
			JP 1670100 C	12-06-1992
			JP 3035347 B	27-05-1991
			JP 57195161 A	30-11-1982
WO 0219836	A	14-03-2002	AU 9155001 A	22-03-2002
			WO 0219836 A2	14-03-2002
			CA 2389244 A1	14-03-2002
			GB 2371732 A	07-08-2002
			US 2003022274 A1	30-01-2003

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

THIS PAGE BLANK (USPTO)